

SOVRANA

ALFALFA

THE QUEEN OF THE QUALITY!





SOVRANA

Type: Winter active (dormancy fall index: 6.5)

Persistency: 4 years and more Protein content (% on D.M): 22.4%

Sowing rate:

- 20 to 25 kg/ha (8 to 10 kg/ac) straight
- 2 Précidose®/ha

Queen of the quality:

- SOVRANA distinguishes itself through its excellent fodder's quality. It has a very high leaf/stem ratio, a very high protein content and is very digestible.
- The fodder is very well consumed and valued by cattle.

Remarkable production:

- SOVRANA shows a very high yielding potential and a great production consistency.
- It's a variety with a good establishment and that distinguishes itself after each cut by a quick and vigorous regrowth.

Hardy:

- SOVRANA has a good level of resistance to all diseases.
- It also has a good ability to resist to dry conditions and to lodging.



is also available with different coating technologies:

ELIFE SAS PREMIUM : mycorrhizae + rhizobium + nutrients cocktail : a technology which optimizes the water and nutrients absorption and improves stress resistance.

GOLD PREMIUM: rhizobium + nutrients cocktail: a ready-to-use product that maximizes the potential of alfalfa.

ENERGY nutrients cocktail: a coating which secures crop establishment and yields while preventing from trace-elements deficiencies.

READY rhizobium: pre-inoculated seeds fo a secured crop development.

Our S.A.S range for alfalfa contains 84% seeds and 16% coating solution, except the S.A.S READY grade with 70% seeds and 30% coating solution. Some S.A.S grades are available in Precidose®, an efficient packing which ensures an accurate number of seeds per bag. Using 2 Precidose®/ha ensures an optimal sowing density of 900 seeds/m².





S.A.S READY . S.A.S GOLD. S.A.S PREMIUM: pre-inoculated seeds, coated with the strain Sinorhizobium meliloti - 2.5/3.10 E8 CFU/o NITRAGIN GOLD - MA n°1150014 - No classification - Proprietary NOVOZYMES A/S Krogshoejvej 36 2880 BAGSVAERD DENMARK. RHIZOSEED - MA n° 1190518 - No classification - Proprietary CYBELE AGROCARE SAS 7 rue Aristide Briand 92 300 LEVALLOIS-PERRET FRANCE.

S.A.S LIFE, S.A.S PREMIUM+: pre-inoculated seeds, coated with the strains Sinorhizobium meliloti - 2.5/3.10 E8 CFU/q and Glomus intraradices 500

propagules/g
NITHAGIN GOLD - MA n°1150014 - No classification - Proprietary NOVOZYMES A/S Krogshoejvej 36 2880 BAGSVAERD DENMARK.
RHIZOSED - MA n° 1190518 - No classification - Proprietary CYBELE AGROCARE SAS 7 rue Aristide Briand 92 300 LEVALLOIS-PERRET FRANCE.
0ZOR® - MA n°1301002 - No classification - Proprietary IF TECH Centre Floriloire 3 rue des Magnolias 49130 LES PONTS DE CE FRANCE.
0ZOR® is a registered trademark of IF TECH.

cérience



FOR WHAT USE?



Sowing:

- Choose suitable fields for cultivating alfalfa: well drained soils and pH above 6.5, consider liming if necessary;
- Ideally sow in spring for a better establishment;
- In summer, drill as soon as the previous crop has been harvested.
- Prepare a thin seed bed to ensure optimal soil-seed contact;
- Sow at a depth of 0.5 to 1 cm maximum. If bare or S.A.S ENERGY coated seeds are used, it is advised to inoculate the seeds with a Rhizobium meliloti strain prior to sowing, particularly if no alfalfa has been cultivated for the past years, but also in case the organic matter content of the soil is low.
- Roll right after sowing.

Protection:

- Monitor slugs during the establishment and take action if necessary.
- Control weeds at 3-4 trifoliate leaves stage if needed.

- Perform soil analysis regularly in order to know the pH and the nutrients contents with accuracy. Adjust fertilization accordingly;
- Make sure you compensate the nutrients that are exported and increase your inputs in case of poor soils.
- Alfalfa is very demanding in potassium ($\rm K_20$) and, to a lower extend, in phosphorus ($\rm P_2O_5$).



Packaging: 15 kg or 25 kg bag or Précidose® (4.5 milions of seeds)



NOV 2022 V1 -·Non-contractual document.

This document is provided for information purposes only. Recommandations may vary according to local soil and climate conditions and local agricultural practices.