



Cerialience
L'AGRONOME SEMENCIER



M-TALENT

FORAGE MIXTURE

**100% LEGUMES TO BE COMBINED WITH A GRASS
TO PRODUCE HIGH-QUALITY FODDER.**





M-TALENT

Profile:

Winter catch crop

Harvesting period:

6 to 9 months (can be harvested in the autumn depending on biomass production, and again in April–May)

Soil types:

- All soil types

Sowing rate (kg/ha): 10 to 20 kg/ha; to be adjusted according to the companion species and the desired quality of the fodder.

Forage mix (%by weight):

- 45% Crimson clover
- 25% NICKEL hairy vetch
- 20% SAVANE hairy vetch
- 10% Balensa clover



ITS ADVANTAGES

- **M-TALENT offers greater harvesting flexibility:**
 - It is possible to harvest the fodder as early as autumn and/or carry out an early cut in spring.
- **Higher-quality fodder:**
 - Two hairy vetches have been selected for this mixture.
 - Their complementary nature ensures a high content of proteins of the mixture:
 - . NICKEL, semi-early hairy vetch;
 - . SAVANE, late hairy vetch.
- **Great flexibility in combination:**
 - It can be combined with a wide variety of species: R.G.I, rye, etc.
 - The sowing rate should be adapted to the companion species and the desired forage quality.



Since mixing the best varieties does not necessarily provide the best forage mixture, this mixture is made of varieties which have been bred and evaluated for their ability to grow in a multi-species mixture.

This mixture secures production as it will adapt to the heterogeneity within the field and to climatic hazards. To fulfill this objectives and provide the best

performances, this mixture is made of a chosen diversity of species and also of a chosen intra-specific diversity (diversity of varieties within individual species).

FOR WHICH SITUATIONS?

Grazing	Mowing

Sowing:

- Prepare soil that is finely crumbled on the surface and firmly tilled deep down;
- Sow at a depth of 1 to 2 cm;
- Roll the soil immediately after sowing.

Protection:

- Monitor for slugs at the time of planting and take action if necessary.

Fertilization:

- Before applying any fertiliser, take into account the nutrients available in the soil by referring to the soil analysis and the manure applied by livestock.
- Apply at planting:
 - 70 to 80 units of Phosphorus/ha;
 - 100 to 150 units of Potash/ha;
- At the end of winter, apply 50 to 80 units of Nitrogen/ha depending on residuals and yield targets.




Packing:
15 kg bag

S.A.S STIM: seeds with a biostimulant to promote planting and enhance stress resistance.
OSYR – Marketing Authorisation No. 108002 – Non-classified product –
Property of FRAYSSINET, La Mothe 81240 ROUAIROUX.
Our S.A.S STIM coated seeds contain 70% bare seeds and 30% coating.

Do not allow hairy vetch to be consumed in seed form.



Route de la Métré - Beaufort-en-Vallée –
49250 BEAUFORT EN ANJOU - FRANCE
Tél : + 33 (0)5 49 54 20 54
www.cerience.fr



March 2026/V1-Non-binding document.
The information contained in this document is provided for guidance only and may vary depending on pedoclimatic conditions and growing conditions.
The varieties included in the mixture are subject to change depending on availability.