



Cérissance
L'AGRONOME SEMENCIER



AYACARA

ANTI-NEMATODE FORAGE RADISH

Multiple resistance to nematodes and clubroot



AYACARA

PROPERTIES

- Species: Fodder radish
- Origin: Vandinter/Cérieence
 - Registration: Germany 2022
 - Flowering: late to very late
 - PMG: 12 to 14 g
 - Sowing rate: 9 to 20 g/ha pure

ACTION

Nematocide

- Heterodera schachtii (beetroot): BCA1
- Meloidogyne chitwoodi (potato, corn): Resistant
- Meloidogyne incognita (corn, vegetables): Resistant
- Cabbage clubroot: Resistant

ADVANTAGES

- Very high nematicidal action against beet nematodes (BCA1).
- Double resistance to potato, maize and vegetable nematodes.
- Resistance to clubroot.
- Very late flowering.
- Very rapid ground cover.


OFFICIAL RESULTS

German catalogue of anti-nematode mustard varieties 2025

Varieties	Years of registration	Resistance to cyst nematodes - <i>H. schachtii</i>	Resistance to the gall nematode <i>M. chitwoodi</i>	Soil coverage speed (1 = poor, 9 = very good)	Late flowering (1 = very late/ 9 = very early)	** Resistance to cabbage hernia
AYACARA	2022	BCA 1	R	7	4	R
Witness A	2018	BCA 2	R	5	3	/
Tajuna	2012	BCA 2	R	6	5	/
Witness C	2015	BCA 2	R	6	4	/
Trident	2015	BCA 2	R	7	3	/
Mercator	2013	BCA 2	/	7	4	/

* Resistance to cabbage hernia = test result from SNES (France)
 BCA 1 = nematode reduction >99%

FOR WHICH SITUATION?

Cover	Companion plant oilseed rape
	

Planting advice:

- Manage crop residues
- Shallow stubble cultivation to a maximum depth of 3 to 5 cm
- Sow at a depth of 1 to 2 cm
- Roll to ensure seed-soil contact



Packing:
Bag de 25 kg